


DTIC FILE COPY
Naval Research Laboratory

Washington, DC 20375-5000



NRL Memorandum Report 6606

AD-A218 146

On Free Products of Torsion Free Abelian Groups

A. M. GAGLIONE

*Identification Systems Branch
Radar Division*

February 23, 1990

DTIC
ELECTE
FEB 22 1990
S D

Approved for public release; distribution unlimited.

90 02 21 020

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release, distribution unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 6606		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b. OFFICE SYMBOL (If applicable) Code 5350	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Air Systems Command	8b. OFFICE SYMBOL (If applicable) APC-209C	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Washington, DC 20361-1213		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 64211N	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO. DN180-248
11. TITLE (Include Security Classification) On Free Products of Torsion Free Abelian Groups			
12. PERSONAL AUTHOR(S) * Gaglione, A.M.			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM Jul 88 TO Jul 89	14. DATE OF REPORT (Year, Month, Day) 1990 February 23	15. PAGE COUNT 16
16. SUPPLEMENTARY NOTATION *Affiliation: Department of Mathematics, U.S. Naval Academy, Annapolis, MD 21402-5000			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) group, free product, commutator, lower central series, basic commutator, F-simple commutator, J-simple commutator, J-basic commutator	
FIELD	GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Let G be a group. For $x, y \in G$ the commutator of x and y is $(x, y) = x^{-1} y^{-1} x y.$ The lower central series of G is a sequence of normal subgroups of G defined inductively as follows: $G_1 = G$, and having defined G_n (for n a positive integer), G_{n+1} is defined as the group generated by all commutators of (x, y) where $x \in G_n$ and $y \in G$. In this paper the lower central series is studied for a group J which is a free product of a finite number of torsion free Abelian groups. In particular, the quotient groups J_n / J_{n+1} are completely determined.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Emanuel Vegh		22b. TELEPHONE (Include Area Code) (202) 767-3481	22c. OFFICE SYMBOL Code 5350

CONTENTS

1. INTRODUCTION	1
2. PRELIMINARIES	1
3. STATEMENT AND PROOF OF THE MAIN RESULT	3
CONCLUSION	12
REFERENCES	12



Accession For	
PTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Date	Avail and/or Special
A-1	

ON FREE PRODUCTS OF TORSION FREE ABELIAN GROUPS

1. INTRODUCTION

In a previous paper [1], the quotient groups of the lower central series $\bar{G}_n = G_n / G_{n+1}$ were studied. There the group G was assumed to be a free product of a finite number of finitely generated Abelian groups and G_n denoted the n^{th} subgroup of the lower central series of G . Here we give an improved proof of a complicated lemma which first appeared in [1] (in particular, Lemma 4.4[1]). The proof given here, especially for property (iii) of the conclusion of that lemma, is a significant simplification of that which appears in [1]. We observe that one of the consequences of Lemma 4.4[1] is to give a set of free generators for the lower central quotients in the case where the free factors are torsion free (i.e., $G = J$ in the terminology of [1]. Moreover the free generators are the J -basic commutators also using the terminology of Definition 4.1 of [1].). The author only thought of this simplification after the publication of [1]. Furthermore, the improved proof uses results from [2] and [3].

In this paper, notation, terminology, results, references, and equations of [1] will be employed. Furthermore, the numbering, a.b, of any definition, equation, etc., of [1] will correspond here to a.b-I. For example, Lemma 2.1-I will mean Lemma 2.1 of [1].

2. PRELIMINARIES

In order to carry out our goal, we need to give some preliminary machinery.

Lemma 2.1. (cf. Corollary 5.1 [3]) Let S be the subgroup of

F generated by a finite set Σ of F -simple commutators (see Definition 3.6-I) of weight > 1 . Let $U = \{u_1, u_2, \dots, u_m\} \subseteq \Sigma$. Let $V = \{v_1, v_2, \dots, v_m\}$ be a set of m nontrivial elements v_i of F such that $W(v_i) \geq W(u_i)$ for $1 \leq i \leq m$. (Here we are using the notation of Definition 2.1-I for weight.) Let ψ be the mapping of Σ given by

$$\begin{aligned} u_i &\longrightarrow v_i \text{ if } u_i \in U \\ c_i &\longrightarrow c_i \text{ if } c_i \notin U. \end{aligned}$$

Then ψ extends to a homomorphism, also denoted by ψ , of S into F such that if $u \in S$ and $W(u) = \omega > 0$, then its image $v = \psi(u) \in F_\omega$.

We also will require

Lemma 2.2. (cf. Theorem 1.1[2]) Let c be the F -simple basic commutator of weight $n > 1$,

$$c = (c_{j_1}, c_{j_2}, \dots, c_{j_n}). \quad (2.1)$$

Let d be a generator of F such that $d < c_{j_n}$. Let $v =$

$(c_{j_2}, d, e_1, e_2, \dots, e_{n-1})$ be such that $e_1 = c_{j_1}$ for $n = 2$, but

$e_1 \leq e_2 \leq \dots \leq e_{n-1}$ is a rearrangement of $c_{j_1}, c_{j_3}, \dots, c_{j_n}$, for

$n > 2$. (Note that v is F -simple of weight $n + 1$ if $d < c_{j_2}$.) Then

the following identity holds in F

$$(c,d) = \left\{ \begin{array}{ll} \pi_1 M(c,d) \pi_2 & \text{if } d \geq c_{j_2} \text{ (Case I)} \\ \pi_1 v^{-1} \pi_2 M(c,d) \pi_3 & \text{if } d < c_{j_2} \text{ (Case II)} \end{array} \right\} \quad (2.2)$$

where the π_j ($j = 1, 2, 3$) are either $= 1$ or are words in finitely many F-simple commutators u_i such that

- (i) $1 < W(u_i) < n + 1$;
- (ii) if $u_i = (v_1, v_2, \dots, v_t)$, then v_1, v_2, \dots, v_t is a rearrangement of a subsequence of $c_{j_1}, \dots, c_{j_n}, d$.

3. STATEMENT AND PROOF OF THE MAIN RESULT

We are now ready to state and prove our main result

Theorem 3.1. (See Lemma 4.4-I.) Let c be F-simple as in (2.1). Suppose that c satisfies Criterion 1 but not Criterion 3. (See Definition 4.1-I.) Let t be the largest integer such that c_{j_t} occurs in c and $(c_{j_t}, c_{j_1}) = 1$ in J . (The hypothesis that c does not satisfy Criterion 3 guarantees that such a t exists, $t \geq 3$, and that $j_t > j_1$.) Let the F-simple commutator $d(c)$ be defined by

$$d(c) = (c_{j_t}, d_2, d_3, \dots, d_n) \quad (3.1)$$

where d_2, d_3, \dots, d_n is that rearrangement of $c_{j_1}, \dots, c_{j_{t-1}}, c_{j_{t+1}}, \dots, c_{j_n}$ for which $d_2 \leq \dots \leq d_n$. Then the following holds in J

$$c = H_1 d(c) H_2 \quad (3.2)$$

where the H_j ($j = 1, 2$) are words in J -simple commutators u_i such that (i) $1 < W(u_i) < n$; (ii) if $u_i = (v_1, \dots, v_k)$, then v_1, \dots, v_k is a rearrangement of a subsequence of c_{j_1}, \dots, c_{j_n} ; (iii) $W(H_1 d(c) H_2) \geq n$ in F .

Proof. We proceed by induction on the place of c in our lexicographic ordering of basic commutators. Clearly, $W(c) = n \geq 3$. For $n = 3$, we have $c = (c_{j_1}, c_{j_2}, c_{j_3})$ with $(c_{j_1}, c_{j_3}) = 1$ in J and $c_{j_3} > c_{j_1}$. Thus equation (3.20)-I yields that in J

$$c = (c_{j_1}, c_{j_2})^{-1} (c_{j_3}, c_{j_2}) (c_{j_3}, c_{j_2}, c_{j_1}) \cdot (c_{j_1}, c_{j_2}) (c_{j_3}, c_{j_2})^{-1}. \quad (3.3)$$

This shows that we have (3.2) with properties (i) and (ii) for $n = 3$. In order to show that the right hand side of (3.3) also satisfies property (iii), we let $a = (c_{j_1}, c_{j_2})$, $b = (c_{j_3}, c_{j_2})$, and $A = (c_{j_3}, c_{j_2}, c_{j_1})$. Then (3.3) becomes

$$c = a^{-1} b A a b^{-1}$$

and we need to show that $W(a^{-1} b A a b^{-1}) \geq 3$ in F . But in F

$$\begin{aligned} a^{-1} b A a b^{-1} &= (a^{-1} b A b^{-1} a) \cdot (a, b^{-1}) \\ &= R \cdot (a, b^{-1}) \end{aligned}$$

where $R = a^{-1} b A b^{-1} a$. Evidently by (2.3a)-I, $W(a, b^{-1}) \geq 4$ in F so that

$$W(a^{-1} b A a b^{-1}) = W(R(a, b^{-1})) \geq W(R) \geq 3$$

in F . This proves the result for $n = W(c) = 3$.

Next, suppose that we have already proven the result for all $c < c_k$ with $W(c_k) > 3$. We then proceed to the smallest commutator c which satisfies the hypothesis of our theorem and is such that $c \geq c_k$. (Thus $n = W(c) \geq W(c_k) > 3$.)

We now define $\delta = n - t$ and we treat two cases: $\delta > 0$ and $\delta = 0$. For $\delta > 0$, let us write $e = c^L$ and we note that, in J , $e = (c_{j_1}, \dots, c_{j_{n-1}})$ is by the induction hypothesis a word of the form (3.2), i.e.,

$$e = H_{11} d(e) H_{12},$$

where H_{1j} ($j = 1, 2$) are words in J -simple basic commutators z_i satisfying (i) and (ii) of our conclusion, and $W(H_{11} d(e) H_{12}) \geq n - 1$ in F . In particular every z_i has $1 < W(z_i) < n - 1$ and also

has $z_i^R \leq c_{j_{n-1}}$ (recall by hypothesis that $c_{j_1} < c_{j_t} \leq c_{j_{n-1}}$).

Applying the trivial identity (3.19)-I to the computation of $c = (e, c_{j_n}) = (H_{11} d(e) H_{12}, c_{j_n})$, we find that in J

$$c = \Pi_1 (d(e), c_{j_n}) \Pi_2 \quad (3.4)$$

where the Π_j ($j = 1, 2$) are words in z_i , (z_i, c_{j_n}) , and $d(e)$. (Note $d(e)$ only occurs in Π_1 .) Now $d(c) = (d(e), c_{j_n})$ by (3.1) since $\delta > 0$. Here we denote the word on the right hand side of (3.4) by

w and note that w is a word in the F-simple basic commutators $d(e)$, z_i , (z_i, c_{j_n}) , and $d(c)$, i.e.,

$$w = \Pi_1 d(c) \Pi_2 \quad (3.4a)$$

where the Π_j ($j = 1, 2$) are as in (3.4). Moreover, we note that $W(z_i, c_{j_n}) \leq n - 1$. So that either the F-simple commutators (z_i, c_{j_n}) are all J-simple or we may use the induction hypothesis to write those (z_i, c_{j_n}) which are not J-simple as words of the form (3.2) in J-simple commutators. After making the replacements for those (z_i, c_{j_n}) in (3.4a) which are not J-simple, we find that in J

$$c = H_1 d(c) H_2 \quad (3.5)$$

where we have demonstrated that the H_j ($j = 1, 2$) are words in J-simple commutators satisfying conditions (i) and (ii) of our conclusion.

To see that our expression for c in (3.5) also satisfies condition (iii) of our conclusion, we first note that the word w as given in (3.4a) has $W(w) \geq n$ in F. (This is true since $w = (H_{11} d(e) H_{12}, c_{j_n})$ in F and $W(H_{11} d(e) H_{12}) \geq n - 1$.) Thus if all the (z_i, c_{j_n}) satisfy Criterion 3, i.e., are J-simple, we are done. Otherwise let Σ be set of all the distinct F-simple basic commutators which occur in w as given by (3.4a). Let $U = (u_1, \dots, u_m) \subseteq \Sigma$ be those (z_i, c_{j_n}) in (3.4a) which are not

J-simple and let $\{v_1, \dots, v_m\}$ be those words in J-simple commutators which are obtained by applying the induction hypothesis to the (z_i, c_{j_n}) . Thus $u_i = v_i$ in J and $W(u_i) \leq W(v_i)$ in F for all $i = 1, \dots, m$. Now to go from (3.4) to (3.5), we replace the u_i by the v_i in w as given by (3.4a), i.e., we apply the map $\psi : u_i \rightarrow v_i$. Thus since $w \in F_n$, Lemma 2.1 implies that

$$\psi(w) = H_1 d(c) H_2$$

in (3.5) is such that $(H_1 d(c) H_2) \in F_n$, i.e., $W(H_1 d(c) H_2) \geq n$.

This completes the proof of our result in the case $\delta > 0$.

It remains to consider the case $\delta = 0$ for which we write

$(c^L)^L = A$, $(c^L)^R = c_{j_n} = g$, and $c^R = h$. Then identity

(3.20)-I gives

$$c = (A, g, h) = (A, g)^{-1} (A, h)^{-1} A^{-1} (h, g)$$

$$A(A, g)(A, h)(A, h, g)(h, g)^{-1}. \quad (3.6)$$

Either $(h, g) = 1$ in J or it is J-simple. Also all the commutators A , (A, g) , and (A, h) are F-simple of weight $< n$. Hence any such commutator is either J-simple or can be written in J by the induction hypothesis as a word in J-simple commutators satisfying conditions (i), (ii), and (iii) of our conclusion. Letting

$$Z = (A, h, g)$$

in (3.6), we claim that is sufficient to prove that Z can be written in J as a word in J-simple commutators in the form

$$Z = H_{11} d(c) H_{12} \quad (3.7)$$

where the H_{1j} ($j = 1, 2$) are words in J -simple commutators, u_i , satisfying (i) and (ii) of our conclusion, and

$$W(H_{11} d(c) H_{12}) \geq n.$$

To see this, suppose we substitute (3.7) and the expressions in terms of J -simple commutators (for those among A , (A, g) , and (A, h) which are not J -simple) into (3.6). This writes c in J as

$$c = H_1 d(c) H_2 \quad (3.8)$$

where the H_j ($j = 1, 2$) are words in J -simple commutators. Clearly, conditions (i) and (ii) of our conclusion are satisfied by (3.8), according to the induction hypothesis and our claim.

For condition (iii), we note that (3.6) is an identity in F . Thus the right hand side of (3.6) has weight n in F since c is F -simple of weight n . Now let $(h, g) = d$, $(A, g) = a$, and $(A, h) = b$. Then (3.6) becomes

$$\begin{aligned} c &= a^{-1} b^{-1} A^{-1} d A a b Z d^{-1} \\ &= (a, b) (A a b, d^{-1}) (d Z d^{-1}). \end{aligned} \quad (3.6a)$$

By (2.3a-I), $a \in F_{n-1}$, $b \in F_{n-1}$, $A \in F_{n-2}$, $Z \in F_n$, and $d \in F_2$.

(Either $d = 1$ or $W(d) = 2$.) Let \tilde{a} , \tilde{b} , \tilde{A} , \tilde{Z} be the words in J -simple commutators such that $a = \tilde{a}$, $b = \tilde{b}$, $Z = \tilde{Z}$ in J (i.e., $\tilde{Z} = H_{11} d(c) H_{12}$ as in (3.7)). We know by the induction hypothesis

and our claim for Z that $\tilde{a} \in F_{n-1}$, $\tilde{b} \in F_{n-1}$, $\tilde{A} \in F_{n-2}$, $\tilde{Z} \in F_n$.

Also $\tilde{d} \in F_2$ since d either is 1 in J or it is a J -simple

commutator of weight 2. (Here $\tilde{d} = d$ if $d \neq 1$ in J but $\tilde{d} = 1$ if $d = 1$ in J .) Furthermore, the mapping of $F \rightarrow J$ is a homomorphism; also the J -simple commutators are free generators of the subgroup of F_2 which they generate by Corollary 4.1-I. Thus

$$\tilde{c} = (\tilde{a}, \tilde{b}) (\tilde{A} \tilde{a} \tilde{b}, \tilde{d}^{-1}) (\tilde{d} \tilde{Z} \tilde{d}^{-1})$$

is according to (2.3a)-I such that $(\tilde{a}, \tilde{b}) \in F_{2n-2} \subseteq F_n$ (since

$2n - 2 > n$ in our case $n > 3$), $(\tilde{A} \tilde{a} \tilde{b}, \tilde{d}^{-1}) \in F_n$, and

$\tilde{d} \tilde{Z} \tilde{d}^{-1} \in F_n$. Thus $W(\tilde{c}) \geq n$ in F where \tilde{c} is the unique word in

J -simple commutators which $= c$ in J . (Here $\tilde{c} = H_1 d(c) H_2$ as in (3.8).)

Thus we now focus our attention on $Z = (A, h, g)$, and prove the claim for Z as given by (3.7). As already mentioned, $b = (A, h)$ is, by the induction hypothesis, a word, \tilde{b} , in J of the form (3.2) in $d(b)$ and J -simple commutators, s_i , of weight $< n-1$ with all the special properties of our conclusion. Therefore $Z = (b, g)$ can be written by the trivial identity (3.19)-I in J in the form

$$Z = (\tilde{b}, g) = H_{01}(d(b), g) H_{02} \quad (3.9)$$

where the H_{0j} ($j = 1, 2$) are words in $d(b)$, s_i , and (s_i, g) . By the

induction hypothesis $W(\tilde{b}) \geq n - 1$, hence (2.3a)-I implies that the right side of (3.9), $H_{01}(d(b), g)H_{02}$, has weight at least n in F .

Let us now examine $(d(b), g)$. From (3.1) and the fact that $\delta = 0$,

$$d(b) = (c_{j_n}, e_2, e_3, \dots, e_{n-1})$$

where $e_2 \leq e_3 \leq \dots \leq e_{n-1}$ is a rearrangement of

$c_{j_1}, c_{j_2}, \dots, c_{j_{n-2}}$. (Evidently $c_{j_2} = e_2$, since $c_{j_1} > c_{j_2} \leq c_{j_3} \leq \dots$

$\leq c_{j_n}$.) We note from Lemma 3.5-I that $M(d(b), g) = d(c)$. Next, we

apply Lemma 2.2 to write $(d(b), g)$ as $\Pi_1 d(c) \Pi_2$ in F . (Note that

$g = c_{j_{n-1}} \geq e_2$ so that we are in Case I of Lemma 2.2 and also that if

$g \geq e_{n-1}$, then $(d(b), g) = d(c)$ and we just then take $\Pi_1 = \Pi_2 =$

1.)

Let us also apply Lemma 2.2 to rewrite those (s_i, g) in (3.9)

which are not basic (i.e., $g < s_i^R$ so that Lemma 2.2 applies) in terms of F -simple commutators. Thus

$$(s_i, g) = \Pi_{i1} M(s_i, g) \Pi_{i2}$$

in F . (Here it does not matter which case of (2.2) we are in

because $W(s_i, g) < n$.) Thus Π_1, Π_2 , and Π_{ij} ($j = 1, 2$) are words

in F -simple commutators, z_i , all of which have $1 < W(z_i) < n$

and satisfy condition (ii) of our conclusion. Substituting these into (3.9), we have that Z can be written in J as a word, w , in

F -simple commutators $d(c), s_i, (s_i, g)$ (only those which are basic

are included here), and z_i , i.e., in J we have

$$Z = w = \Omega_{01} d(c) \Omega_{02} \quad (3.10)$$

where Ω_{0j} ($j = 1, 2$) are words in the s_i , (s_i, g) -- only the basic ones -- , and z_i . We may now use the induction hypothesis to write those F -simple commutators, the (s_i, g) and z_i , which occur in (3.10) and are not J -simple, as words in J -simple commutators. Making these replacements in (3.10) gives (3.7). To see this, we note that now Z is written, in J , in terms of J -simple commutators, u_i , which clearly satisfy (i) and (ii) of our conclusion.

To see that $W (H_{11} d(c) H_{12}) \geq n$ in F , we first note that the right hand side of (3.9) is equal to the right hand side of (3.10) in F , i.e.,

$$w = H_{01} (d(b), g) H_{02}$$

holds in F . Thus $w \in F_n$. We can now continue in a manner exactly analogous to that used in the case of $\delta > 0$, because the situation in (3.10) is similar to that of equation (3.4a). In this way Lemma 2.1 shows that the weight of $H_{11} d(c) H_{12}$ given in (3.7) is at least n . This completes the proof of our claim for Z given in (3.7) and with that the proof of our theorem is accomplished.

CONCLUSION

Theorem 3.1 of this paper shows that in a group which is a free product of torsion free Abelian groups, here denoted by J , any F -simple commutator can be expressed in terms of J -simple commutators. It then follows from well known results [1] that the J -basic commutators of weight n form a basis for the quotient group J_n / J_{n+1} . Thus Theorem 3.1 completely determines these quotients.

REFERENCES

1. A.M. Gaglione and H.V. Waldinger, Factor groups of the lower central series of free products of finitely generated Abelian groups, *Proc. Groups St. Andrews--1985*, Cambridge University Press, 164-203.
2. _____, A theorem in the commutator calculus, *Proc. Singapore Group Theory Conference 1987*, Walter de Gruyter, 367-371.
3. _____, The commutator collection process, to appear in *Proc. AMS Special Session on Combinatorial Group Theory*, (1988), Amer. Math. Soc.